

**Astellas Submits Supplemental New Drug Application  
in Japan for PADCEV™ (enfortumab vedotin  
(genetical recombination)) with KEYTRUDA®  
(pembrolizumab (genetical recombination)) for First-  
Line Treatment of Advanced Bladder Cancer**

*– Pivotal trial found the enfortumab vedotin plus pembrolizumab combination significantly extended overall and progression-free survival vs platinum-containing chemotherapy–*

*– If approved, PADCEV with KEYTRUDA would be the first combination in Japan to offer an alternative to chemotherapy, the current standard of care in first-line locally advanced or metastatic urothelial cancer –*

**TOKYO – January 31, 2024** – Astellas Pharma Inc. (TSE:4503, President and CEO: Naoki Okamura, “Astellas”) today announced the submission of a Supplemental New Drug Application (sNDA) on January 31 to Japan's Ministry of Health, Labour and Welfare (MHLW) for PADCEV™ (enfortumab vedotin (genetical recombination)) with KEYTRUDA® (pembrolizumab (genetical recombination)) as a combination therapy for the first-line treatment of adult patients with previously untreated locally advanced or metastatic urothelial cancer (la/mUC). If approved, PADCEV with KEYTRUDA has the potential to change the treatment paradigm, becoming the first combination treatment to offer an alternative to platinum-containing chemotherapy, the current standard of care in first-line la/mUC.

Globally, approximately 573,000 new cases of bladder cancer and 212,000 deaths are reported annually.<sup>i</sup> It is estimated that approximately 25,000 people in Japan are diagnosed with bladder cancer each year and approximately 10,000 deaths were reported in 2022.<sup>ii</sup>

**Ahsan Arozullah, M.D., M.P.H., Senior Vice President, Head of Oncology Development, Astellas**

“The initiation of MHLW’s review of our application for enfortumab vedotin and pembrolizumab is encouraging, as we are working to improve upon current treatment options for patients in Japan with advanced stage urothelial cancer, who face poor outcomes at the advanced stage. This submission brings us one step closer to the possibility of offering these patients a treatment that demonstrated improved survival and slowed disease progression compared to platinum-containing chemotherapy.”

The sNDA for the first-line use of this combination is based on results from the Phase 3 EV-302 clinical trial (also known as KEYNOTE-A39). The study found the

combination improved overall survival (OS) and progression-free survival (PFS) with statistically significant and clinically meaningful results in patients with previously untreated la/mUC compared to platinum-containing chemotherapy. The safety results were consistent with those previously reported with this combination, and no new safety issues were identified.

The Committee for Medicinal Products for Human Use (CHMP) of the European Medicines Agency is also reviewing the combination therapy. The U.S. Food and Drug Administration approved the combination therapy in December 2023.

### **About EV-302**

The EV-302 trial is an open-label, randomized, controlled Phase 3 study, evaluating enfortumab vedotin in combination with pembrolizumab versus platinum-containing chemotherapy in patients with previously untreated la/mUC. The study enrolled 886 patients with previously untreated la/mUC who were eligible for cisplatin- or carboplatin-containing chemotherapy regardless of PD-L1 status. Patients were randomized to receive either enfortumab vedotin in combination with pembrolizumab or platinum-containing chemotherapy. The dual primary endpoints of this trial are OS and PFS per RECIST v1.1 by blinded independent central review (BICR). Select secondary endpoints include ORR per RECIST v1.1 by BICR, DOR per RECIST v1.1 by BICR, and safety.

The EV-302 trial is part of an extensive clinical program evaluating this combination in multiple stages of urothelial cancer and other solid tumors. Findings from EV-302 were presented at the European Society for Medical Oncology (ESMO) Congress 2023 in October 2023.

### **About Bladder and Urothelial Cancer**

- Urothelial cancer, or bladder cancer, begins in the urothelial cells, which line the urethra, bladder, ureters, renal pelvis, and some other organs.<sup>iii</sup>
- If bladder cancer has spread to surrounding organs or muscles, it is called locally advanced disease. If the cancer has spread to other parts of the body, it is called metastatic disease.<sup>iv</sup>
- Urothelial cancer accounts for 90% of all bladder cancers and can also be found in the renal pelvis, ureter, and urethra.<sup>iii</sup>
- Approximately 12% of cases are locally advanced or metastatic urothelial cancer at diagnosis.<sup>v</sup>

### **Ongoing Investigational Trials**

The EV-302 trial ([NCT04223856](#)) is an open-label, randomized, controlled Phase 3 study, evaluating enfortumab vedotin in combination with pembrolizumab versus chemotherapy in patients with previously untreated locally advanced or metastatic urothelial cancer (la/mUC) who were eligible for cisplatin- or carboplatin-containing chemotherapy regardless of PD-L1 status.

The EV-103 trial ([NCT03288545](#)) is an ongoing, multi-cohort, open-label, multicenter Phase 1b/2 study investigating enfortumab vedotin alone or in combination with pembrolizumab and/or chemotherapy in first- or second-line settings in patients with la/mUC and in patients with muscle-invasive bladder cancer (MIBC).

Enfortumab vedotin in combination with pembrolizumab is being investigated in an extensive program in multiple stages of urothelial cancer, including two Phase 3 clinical trials in MIBC in EV-304 ([NCT04700124](#), also known as KEYNOTE-B15) and EV-303 ([NCT03924895](#), also known as KEYNOTE-905). The use of enfortumab vedotin in combination with pembrolizumab in second-line urothelial cancer and in MIBC has not been proven safe or effective.

The EV-202 trial ([NCT04225117](#)) is an ongoing, multi-cohort, open-label, multicenter Phase 2 study investigating enfortumab vedotin alone in patients with previously treated advanced solid tumors. This study also has a cohort that is investigating enfortumab vedotin in combination with pembrolizumab in patients with previously untreated recurrent/ metastatic head and neck squamous cell carcinoma.

### **About PADCEV™ (enfortumab vedotin (genetical recombination))**

PADCEV (enfortumab vedotin (genetical recombination)) is a first-in-class antibody-drug conjugate (ADC) that is directed against Nectin-4, a protein located on the surface of cells and highly expressed in bladder cancer.<sup>vi</sup> Nonclinical data suggest the anticancer activity of PADCEV is due to its binding to Nectin-4-expressing cells, followed by the internalization and release of the anti-tumor agent monomethyl auristatin E (MMAE) into the cell, which result in the cell not reproducing (cell cycle arrest) and in programmed cell death (apoptosis).<sup>vii</sup>

### **PADCEV (enfortumab vedotin-ejfv) U.S. Indication & Important Safety Information**

#### **BOXED WARNING: SERIOUS SKIN REACTIONS**

- PADCEV can cause severe and fatal cutaneous adverse reactions including Stevens-Johnson syndrome (SJS) and Toxic Epidermal Necrolysis (TEN), which occurred predominantly during the first cycle of treatment, but may occur later.
- Closely monitor patients for skin reactions.
- Immediately withhold PADCEV and consider referral for specialized care for suspected SJS or TEN or severe skin reactions.
- Permanently discontinue PADCEV in patients with confirmed SJS or TEN; or Grade 4 or recurrent Grade 3 skin reactions.

#### **Indication**

PADCEV®, in combination with pembrolizumab, is indicated for the treatment of adult patients with locally advanced or metastatic urothelial cancer (mUC).

PADCEV, as a single agent, is indicated for the treatment of adult patients with locally advanced or mUC who:

- have previously received a programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PD-L1) inhibitor and platinum-containing chemotherapy, or
- are ineligible for cisplatin-containing chemotherapy and have previously received one or more prior lines of therapy.

### **IMPORTANT SAFETY INFORMATION**

#### **Warnings and Precautions**

**Skin reactions** Severe cutaneous adverse reactions, including fatal cases of SJS or TEN occurred in patients treated with PADCEV. SJS and TEN occurred predominantly during the first cycle of treatment but may occur later. Skin reactions occurred in 70% (all grades) of the 564 patients treated with PADCEV in combination with pembrolizumab in clinical trials. When PADCEV was given in combination with pembrolizumab, the incidence of skin reactions, including severe events, occurred at a higher rate compared to PADCEV as a single agent. The majority of the skin reactions that occurred with combination therapy included maculo-papular rash, macular rash and papular rash. Grade 3-4 skin reactions occurred in 17% of patients (Grade 3: 16%, Grade 4: 1%), including maculo-papular rash, bullous dermatitis, dermatitis, exfoliative dermatitis, pemphigoid, rash, erythematous rash, macular rash, and papular rash. A fatal reaction of bullous dermatitis occurred in one patient (0.2%). The median time to onset of severe skin reactions was 1.7 months (range: 0.1 to 17.2 months). Skin reactions led to discontinuation of PADCEV in 6% of patients.

Skin reactions occurred in 58% (all grades) of the 720 patients treated with PADCEV as a single agent in clinical trials. Twenty-three percent (23%) of patients had maculo-papular rash and 34% had pruritus. Grade 3-4 skin reactions occurred in 14% of patients, including maculo-papular rash, erythematous rash, rash or drug eruption, symmetrical drug-related intertriginous and flexural exanthema (SDRIFE), bullous dermatitis, exfoliative dermatitis, and palmar-plantar erythrodysesthesia. The median time to onset of severe skin reactions was 0.6 months (range: 0.1 to 8 months). Among patients experiencing a skin reaction leading to dose interruption who then restarted PADCEV (n=75), 24% of patients restarting at the same dose and 24% of patients restarting at a reduced dose experienced recurrent severe skin reactions. Skin reactions led to discontinuation of PADCEV in 3.1% of patients.

Monitor patients closely throughout treatment for skin reactions. Consider topical corticosteroids and antihistamines, as clinically indicated. For persistent or recurrent Grade 2 skin reactions, consider withholding PADCEV until Grade  $\leq$ 1. Withhold PADCEV and refer for specialized care for suspected SJS, TEN or for Grade 3 skin reactions. Permanently discontinue PADCEV in patients with confirmed SJS or TEN; or Grade 4 or recurrent Grade 3 skin reactions.

**Hyperglycemia and diabetic ketoacidosis (DKA)**, including fatal events, occurred in patients with and without pre-existing diabetes mellitus, treated with PADCEV. Patients with baseline hemoglobin A1C  $\geq$ 8% were excluded from clinical trials. In clinical trials of PADCEV as a single agent, 17% of the 720 patients treated with PADCEV developed hyperglycemia of any grade; 7% of patients developed Grade 3-4 hyperglycemia (Grade 3: 6.5%, Grade 4: 0.6%). Fatal events of hyperglycemia and DKA occurred in one patient each (0.1%). The incidence of Grade 3-4 hyperglycemia increased consistently in patients with higher body mass index and in patients with higher baseline A1C. The median time to onset of hyperglycemia was 0.5 months (range: 0 to 20 months). Hyperglycemia led to discontinuation of PADCEV in 0.7% of patients. Five percent (5%) of patients required initiation of insulin therapy for treatment of hyperglycemia. Of the patients who initiated insulin therapy for treatment of hyperglycemia, 66% (23/35) discontinued insulin at the time of last evaluation.

Closely monitor blood glucose levels in patients with, or at risk for, diabetes mellitus or hyperglycemia. If blood glucose is elevated (>250 mg/dL), withhold PADCEV.

**Pneumonitis/Interstitial Lung Disease (ILD)** Severe, life-threatening or fatal pneumonitis/ILD occurred in patients treated with PADCEV. When PADCEV was given in combination with pembrolizumab, 10% of the 564 patients treated with combination therapy had pneumonitis/ILD of any grade and 4% had Grade 3-4. A fatal event of pneumonitis/ILD occurred in two patients (0.4%). The incidence of pneumonitis/ILD, including severe events, occurred at a higher rate when PADCEV was given in combination with pembrolizumab compared to PADCEV as a single agent. The median time to onset of any grade pneumonitis/ILD was 4 months (range: 0.3 to 26 months).

In clinical trials of PADCEV as a single agent, 3% of the 720 patients treated with PADCEV had pneumonitis/ILD of any grade and 0.8% had Grade 3-4. The median time to onset of any grade pneumonitis/ILD was 2.9 months (range: 0.6 to 6 months).

Monitor patients for signs and symptoms indicative of pneumonitis/ILD such as hypoxia, cough, dyspnea or interstitial infiltrates on radiologic exams. Evaluate and exclude infectious, neoplastic and other causes for such signs and symptoms through appropriate investigations. Withhold PADCEV for patients who develop Grade 2 pneumonitis/ILD and consider dose reduction. Permanently discontinue PADCEV in all patients with Grade 3 or 4 pneumonitis/ILD.

**Peripheral neuropathy (PN)** When PADCEV was given in combination with pembrolizumab, 67% of the 564 patients treated with combination therapy had PN of any grade, 36% had Grade 2 neuropathy, and 7% had Grade 3 neuropathy. The incidence of PN occurred at a higher rate when PADCEV was given in combination with pembrolizumab compared to PADCEV as a single agent. The median time to onset of Grade  $\geq 2$  PN was 6 months (range: 0.3 to 25 months).

PN occurred in 53% of the 720 patients treated with PADCEV as a single agent in clinical trials including 38% with sensory neuropathy, 8% with muscular weakness and 7% with motor neuropathy. Thirty percent of patients experienced Grade 2 reactions and 5% experienced Grade 3-4 reactions. PN occurred in patients treated with PADCEV with or without preexisting PN. The median time to onset of Grade  $\geq 2$  PN was 4.9 months (range: 0.1 to 20 months). Neuropathy led to treatment discontinuation in 6% of patients.

Monitor patients for symptoms of new or worsening PN and consider dose interruption or dose reduction of PADCEV when PN occurs. Permanently discontinue PADCEV in patients who develop Grade  $\geq 3$  PN.

**Ocular disorders** were reported in 40% of the 384 patients treated with PADCEV as a single agent in clinical trials in which ophthalmologic exams were scheduled. The majority of these events involved the cornea and included events associated with dry eye such as keratitis, blurred vision, increased lacrimation, conjunctivitis, limbal stem cell deficiency, and keratopathy. Dry eye symptoms occurred in 30% of patients, and blurred vision occurred in 10% of patients, during treatment with PADCEV. The median time to onset to symptomatic ocular disorder was 1.7 months (range: 0 to 30.6 months). Monitor patients for ocular disorders. Consider artificial tears for



prophylaxis of dry eyes and ophthalmologic evaluation if ocular symptoms occur or do not resolve. Consider treatment with ophthalmic topical steroids, if indicated after an ophthalmic exam. Consider dose interruption or dose reduction of PADCEV for symptomatic ocular disorders.

**Infusion site extravasation** Skin and soft tissue reactions secondary to extravasation have been observed after administration of PADCEV. Of the 720 patients treated with PADCEV as a single agent in clinical trials, 1% of patients experienced skin and soft tissue reactions, including 0.3% who experienced Grade 3-4 reactions. Reactions may be delayed. Erythema, swelling, increased temperature, and pain worsened until 2-7 days after extravasation and resolved within 1-4 weeks of peak. Two patients (0.3%) developed extravasation reactions with secondary cellulitis, bullae, or exfoliation. Ensure adequate venous access prior to starting PADCEV and monitor for possible extravasation during administration. If extravasation occurs, stop the infusion and monitor for adverse reactions.

**Embryo-fetal toxicity** PADCEV can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risk to the fetus. Advise female patients of reproductive potential to use effective contraception during PADCEV treatment and for 2 months after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with PADCEV and for 4 months after the last dose.

## **ADVERSE REACTIONS**

**Most common adverse reactions, including laboratory abnormalities (≥20%) (PADCEV in combination with pembrolizumab)** Increased aspartate aminotransferase (AST), increased creatinine, rash, increased glucose, PN, increased lipase, decreased lymphocytes, increased alanine aminotransferase (ALT), decreased hemoglobin, fatigue, decreased sodium, decreased phosphate, decreased albumin, pruritus, diarrhea, alopecia, decreased weight, decreased appetite, increased urate, decreased neutrophils, decreased potassium, dry eye, nausea, constipation, increased potassium, dysgeusia, urinary tract infection and decreased platelets.

**Most common adverse reactions, including laboratory abnormalities (≥20%) (PADCEV monotherapy)** Increased glucose, increased AST, decreased lymphocytes, increased creatinine, rash, fatigue, PN, decreased albumin, decreased hemoglobin, alopecia, decreased appetite, decreased neutrophils, decreased sodium, increased ALT, decreased phosphate, diarrhea, nausea, pruritus, increased urate, dry eye, dysgeusia, constipation, increased lipase, decreased weight, decreased platelets, abdominal pain, dry skin.

### **EV-302 Study: 440 patients with previously untreated Ia/mUC (PADCEV in combination with pembrolizumab)**

**Serious adverse reactions** occurred in 50% of patients treated with PADCEV in combination with pembrolizumab. The most common serious adverse reactions (≥2%) were rash (6%), acute kidney injury (5%), pneumonitis/ILD (4.5%), urinary tract infection (3.6%), diarrhea (3.2%), pneumonia (2.3%), pyrexia (2%), and

hyperglycemia (2%). **Fatal adverse reactions** occurred in 3.9% of patients treated with PADCEV in combination with pembrolizumab including acute respiratory failure (0.7%), pneumonia (0.5%), and pneumonitis/ILD (0.2%).

Adverse reactions leading to discontinuation of PADCEV occurred in 35% of patients. The **most common adverse reactions (≥2%) leading to discontinuation** of PADCEV were PN (15%), rash (4.1%) and pneumonitis/ILD (2.3%). Adverse reactions leading to dose interruption of PADCEV occurred in 73% of patients. The **most common adverse reactions (≥2%) leading to dose interruption** of PADCEV were PN (22%), rash (16%), COVID-19 (10%), diarrhea (5%), pneumonitis/ILD (4.8%), fatigue (3.9%), hyperglycemia (3.6%), increased ALT (3%) and pruritus (2.5%). Adverse reactions leading to dose reduction of PADCEV occurred in 42% of patients. The **most common adverse reactions (≥2%) leading to dose reduction** of PADCEV were rash (16%), PN (13%) and fatigue (2.7%).

**EV-103 Study: 121 patients with previously untreated Ia/mUC who were not eligible for cisplatin-containing chemotherapy (PADCEV in combination with pembrolizumab)**

**Serious adverse reactions** occurred in 50% of patients treated with PADCEV in combination with pembrolizumab; the most common (≥2%) were acute kidney injury (7%), urinary tract infection (7%), urosepsis (5%), sepsis (3.3%), pneumonia (3.3%), hematuria (3.3%), pneumonitis/ILD (3.3%), urinary retention (2.5%), diarrhea (2.5%), myasthenia gravis (2.5%), myositis (2.5%), anemia (2.5%), and hypotension (2.5%). **Fatal adverse reactions** occurred in 5% of patients treated with PADCEV in combination with pembrolizumab, including sepsis (1.6%), bullous dermatitis (0.8%), myasthenia gravis (0.8%), and pneumonitis/ILD (0.8%). **Adverse reactions leading to discontinuation** of PADCEV occurred in 36% of patients; the most common (≥2%) were PN (20%) and rash (6%). **Adverse reactions leading to dose interruption** of PADCEV occurred in 69% of patients; the most common (≥2%) were PN (18%), rash (12%), increased lipase (6%), pneumonitis/ILD (6%), diarrhea (4.1%), acute kidney injury (3.3%), increased ALT (3.3%), fatigue (3.3%), neutropenia (3.3%), urinary tract infection (3.3%), increased amylase (2.5%), anemia (2.5%), COVID-19 (2.5%), hyperglycemia (2.5%), and hypotension (2.5%). **Adverse reactions leading to dose reduction** of PADCEV occurred in 45% of patients; the most common (≥2%) were PN (17%), rash (12%), fatigue (5%), neutropenia (5%), and diarrhea (4.1%).

**EV-301 Study: 296 patients previously treated with a PD-1/L1 inhibitor and platinum-based chemotherapy (PADCEV monotherapy)**

**Serious adverse reactions** occurred in 47% of patients treated with PADCEV; the most common (≥2%) were urinary tract infection, acute kidney injury (7% each), and pneumonia (5%). **Fatal adverse reactions** occurred in 3% of patients, including multiorgan dysfunction (1%), hepatic dysfunction, septic shock, hyperglycemia, pneumonitis/ILD, and pelvic abscess (0.3% each). **Adverse reactions leading to discontinuation** occurred in 17% of patients; the most common (≥2%) were PN (5%) and rash (4%). **Adverse reactions leading to dose interruption** occurred in 61% of patients; the most common (≥4%) were PN (23%), rash (11%), and fatigue (9%). **Adverse reactions leading to dose reduction** occurred in 34% of patients;

the most common ( $\geq 2\%$ ) were PN (10%), rash (8%), decreased appetite, and fatigue (3% each).

**EV-201, Cohort 2 Study: 89 patients previously treated with a PD-1/L1 inhibitor and not eligible for cisplatin-based chemotherapy (PADCEV monotherapy)**

**Serious adverse reactions** occurred in 39% of patients treated with PADCEV; the most common ( $\geq 3\%$ ) were pneumonia, sepsis, and diarrhea (5% each). **Fatal adverse reactions** occurred in 8% of patients, including acute kidney injury (2.2%), metabolic acidosis, sepsis, multiorgan dysfunction, pneumonia, and pneumonitis/ILD (1.1% each). **Adverse reactions leading to discontinuation** occurred in 20% of patients; the most common ( $\geq 2\%$ ) was PN (7%). **Adverse reactions leading to dose interruption** occurred in 60% of patients; the most common ( $\geq 3\%$ ) were PN (19%), rash (9%), fatigue (8%), diarrhea (5%), increased AST, and hyperglycemia (3% each). **Adverse reactions leading to dose reduction** occurred in 49% of patients; the most common ( $\geq 3\%$ ) were PN (19%), rash (11%), and fatigue (7%).

## DRUG INTERACTIONS

### Effects of other drugs on PADCEV (*Dual P-gp and Strong CYP3A4 Inhibitors*)

Concomitant use with dual P-gp and strong CYP3A4 inhibitors may increase unconjugated monomethyl auristatin E exposure, which may increase the incidence or severity of PADCEV toxicities. Closely monitor patients for signs of toxicity when PADCEV is given concomitantly with dual P-gp and strong CYP3A4 inhibitors.

## SPECIFIC POPULATIONS

**Lactation** Advise lactating women not to breastfeed during treatment with PADCEV and for 3 weeks after the last dose.

**Hepatic impairment** Avoid the use of PADCEV in patients with moderate or severe hepatic impairment.

**For more information, please see the U.S. full Prescribing Information including BOXED WARNING for PADCEV [here](#).**

### About Astellas

Astellas Pharma Inc. is a pharmaceutical company conducting business in more than 70 countries around the world. We are promoting the Focus Area Approach that is designed to identify opportunities for the continuous creation of new drugs to address diseases with high unmet medical needs by focusing on Biology and Modality. Furthermore, we are also looking beyond our foundational Rx focus to create Rx+<sup>®</sup> healthcare solutions that combine our expertise and knowledge with cutting-edge technology in different fields of external partners. Through these efforts, Astellas stands on the forefront of healthcare change to turn innovative science into VALUE for patients. For more information, please visit our website at <https://www.astellas.com/en>.

### About the Pfizer, Astellas and Merck Collaboration



Seagen and Astellas entered a clinical collaboration agreement with Merck to evaluate the combination of Seagen's and Astellas' PADCEV™ (enfortumab vedotin) and Merck's KEYTRUDA® (pembrolizumab) in patients with previously untreated metastatic urothelial cancer. As previously announced, Pfizer Inc. successfully completed its acquisition of Seagen on December 14, 2023. KEYTRUDA is a registered trademark of Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Rahway, NJ, USA (known as MSD outside of the United States and Canada).

### **Astellas Cautionary Notes**

In this press release, statements made with respect to current plans, estimates, strategies and beliefs and other statements that are not historical facts are forward-looking statements about the future performance of Astellas. These statements are based on management's current assumptions and beliefs in light of the information currently available to it and involve known and unknown risks and uncertainties. A number of factors could cause actual results to differ materially from those discussed in the forward-looking statements. Such factors include, but are not limited to: (i) changes in general economic conditions and in laws and regulations, relating to pharmaceutical markets, (ii) currency exchange rate fluctuations, (iii) delays in new product launches, (iv) the inability of Astellas to market existing and new products effectively, (v) the inability of Astellas to continue to effectively research and develop products accepted by customers in highly competitive markets, and (vi) infringements of Astellas' intellectual property rights by third parties.

Information about pharmaceutical products (including products currently in development) which is included in this press release is not intended to constitute an advertisement or medical advice.

###

---

### **Astellas Contacts:**

#### *For Media*

Elysia Wood

(703) 722-4656

[Elysia.Wood@astellas.com](mailto:Elysia.Wood@astellas.com)

#### *For Investors*

Astellas Pharma Inc.

Corporate Communications

+81-3-3244-3202

---

<sup>i</sup> International Agency for Research on Cancer. Cancer Today: bladder globocan 2020 fact sheet (12-2020). <https://gco.iarc.fr/today/data/factsheets/cancers/30-Bladder-fact-sheet.pdf>.

<sup>ii</sup> Cancer Information Service, Cancer Statistics in Japan. Published 2022. [https://ganjoho.jp/public/qa\\_links/report/statistics/pdf/cancer\\_statistics\\_2023\\_data\\_E.pdf](https://ganjoho.jp/public/qa_links/report/statistics/pdf/cancer_statistics_2023_data_E.pdf)

- 
- iii National Cancer Institute. What is bladder cancer?  
<https://www.cancer.gov/types/bladder#:~:text=Types%20of%20bladder%20cancer,bladder%20cancers%20are%20urothelial%20carcinomas>.
- iv American Society of Clinical Oncology. Bladder cancer: introduction (12-2021).  
<https://www.cancer.net/cancer-types/bladder-cancer/introduction>.
- v National Cancer Institute. Cancer stat facts: bladder cancer.  
<https://seer.cancer.gov/statfacts/html/urinb.html>.
- vi Challita-Eid PM, Satpayev D, Yang P, et al. Enfortumab vedotin antibody-drug conjugate targeting nectin-4 is a highly potent therapeutic agent in multiple preclinical cancer models. *Cancer Res* 2016;76(10):3003-13.
- vii PADCEV [package insert]. Northbrook, IL: Astellas Pharma US, Inc.